

THE MEAN CURVATURE EQUATION ON SEMIDIRECT PRODUCTS $\mathbb{R}^2 \rtimes_A \mathbb{R}$: HEIGHT ESTIMATES AND SCHERK-LIKE GRAPHS

ÁLVARO KRÜGER RAMOS

(October 16, 2015)

Abstract

In the ambient space of a semidirect product $\mathbb{R}^2 \rtimes_A \mathbb{R}$, we consider a connected domain $\Omega \subseteq \mathbb{R}^2 \rtimes_A \{0\}$. Given a function $u: \Omega \rightarrow \mathbb{R}$, its π -graph is $\text{graph}(u) = \{(x, y, u(x, y)) \mid (x, y, 0) \in \Omega\}$. In this paper we study the partial differential equation that u must satisfy so $\text{graph}(u)$ has prescribed mean curvature H . Using techniques from quasilinear elliptic equations we prove that if a π -graph has nonnegative mean curvature function, then it satisfies some uniform height estimates that depends on Ω and on the supremum the function attains on the boundary of Ω . When $\text{trace}(A) > 0$, we prove that the oscillation of a minimal graph assuming the same constant value n along the boundary tends to zero when $n \rightarrow +\infty$ and goes to $+\infty$ if $n \rightarrow -\infty$. Furthermore, we use these estimates, allied with techniques from Killing graphs, to prove the existence of minimal π -graphs assuming the value 0 along a piecewise smooth curve γ with endpoints p_1, p_2 and having as boundary $\gamma \cup (\{p_1\} \times [0, +\infty)) \cup (\{p_2\} \times [0, +\infty))$.

Mathematics Subject Classification: Primary 53A10, Secondary 49Q05, 35J62.

Keywords and phrases: Minimal surfaces, metric Lie groups, semidirect products, quasilinear elliptic operator, height estimates, Scherk-like graphs..

1. Introduction

Let $A \in M_2(\mathbb{R})$ be a 2×2 matrix. The semidirect product $\mathbb{R}^2 \rtimes_A \mathbb{R}$ is, as a set, the euclidean 3-space $\mathbb{R}^3$, but endowed with a group operation and with a left invariant metric that come from the exponential map $z \mapsto e^{Az}$. More details about its construction are given in Section 2 below. Also, the work of W. Meeks and J. Pérez [MP] is a good reference on the subject, providing the basic aspects of the geometry in these spaces.

There are two main difficulties when dealing with minimal π -graphs in semidirect products $\mathbb{R}^2 \rtimes_A \mathbb{R}$: the first one is that vertical translations

$(x, y, z) \mapsto (x, y, z + t)$ are not isometries of the ambient space. In particular this affects the mean curvature operator so that the coefficients of its second order terms depend on the solution, and the comparison principle (for instance Theorem 10.1 of [GT] and its generalizations) does not apply. The second one is that, unless $\text{trace}(A) = 0$, constant functions do not provide minimal graphs, so there is no maximum principle, in the sense that the supremum (or infimum) of a solution to the minimal graph equation may be strict and attained in the interior of the domain.

In this paper, we consider a convex domain $\Omega \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ with piecewise smooth boundary and exhibit the partial differential equation a function $u: \Omega \rightarrow \mathbb{R}$ must satisfy for its π -graph

$$\text{graph}(u) = \{(x, y, u(x, y)) \in \mathbb{R}^2 \rtimes_A \mathbb{R} \mid (x, y, 0) \in \Omega\}$$

to have prescribed mean curvature function. Depending on the trace and on the determinant of A such PDE has different behaviours. For instance, when $\text{trace}(A) = 0$, if u is such that $\text{graph}(u)$ has nonnegative mean curvature $H \geq 0$ with respect to the upwards orientation, then it satisfies the maximum principle

$$\sup_{\partial\Omega} u = \sup_{\Omega} u. \tag{1}$$

This property was first observed by W. Meeks, P. Mira, J. Pérez and A. Ros in [MMPR3] (we state this result as Lemma 3.1 below), and we remark that (1) does not hold when $\text{trace}(A) > 0$, even for $H \equiv 0$: a minimal graph that is constant along its boundary necessarily assumes an interior maximum and it is not constant, as horizontal planes (representing constant functions) are no longer minimal. It becomes a natural question to ask if there is a *maximal oscillation* these minimal graphs that are constant along the boundary can attain, and this question is answered in this paper via height estimates of partial differential equations.

Let us describe some of the main results of this paper: in Section 3, given $\Omega \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ and a parameter $\alpha \in \mathbb{R}$, we obtain, in Theorem 3.2, a constant $C(\alpha) = C(\text{diam}(\Omega), \alpha)$ such that if $u: \Omega \rightarrow \mathbb{R}$ is a function such that $\text{graph}(u)$ has nonnegative mean curvature, then

$$\sup_{\partial\Omega} u \leq \alpha \Rightarrow \sup_{\Omega} u \leq \sup_{\partial\Omega} u + C(\alpha). \tag{2}$$

Still in Section 3 we prove that the dependence of α in (2) is essential (Theorem 3.3) for the validity of the result, on the sense that it is not possible to obtain some constant $C = C(\Omega)$ such that every $u: \Omega \rightarrow \mathbb{R}$ such that $\text{graph}(u)$ has nonnegative mean curvature satisfies the uniform height estimate

$$\sup_{\Omega} u \leq \sup_{\partial\Omega} u + C.$$

We also use, in Theorem 3.5, the freedom of the parameter α to obtain that, when $\text{trace}(A) > 0$ the oscillation of a family of solutions to the problem

$$\begin{cases} \text{graph}(u) \text{ is a minimal surface of } \mathbb{R}^2 \rtimes_A \mathbb{R} \\ u|_{\partial\Omega} = \alpha \in \mathbb{R} \end{cases}$$

converges to zero when α approaches $+\infty$. Moreover, we prove it goes to $+\infty$, if $\alpha \rightarrow -\infty$.

We finish the paper in Section 4, where we bring techniques from Killing graphs, in addition to the estimates on the coefficients of the mean curvature operator obtained on Section 3, to generalize an argument of A. Menezes [Me] to any semidirect product $\mathbb{R}^2 \rtimes_A \mathbb{R}$, and obtain the existence of minimal π -graphs which are similar to the fundamental piece of the doubly periodic Scherk surface of $\mathbb{R}^3$, on Theorem 4.1.

2. Semidirect products $\mathbb{R}^2 \rtimes_A \mathbb{R}$

This section is to give a brief review about semidirect products $\mathbb{R}^2 \rtimes_A \mathbb{R}$. We follow the notation and construction of W. Meeks and J. Pérez, [MP].

Let H, V to be two groups and let $\varphi: V \rightarrow \text{Aut}(H)$ a group homomorphism between V and the group of automorphisms of H . Then, the *semidirect product between H and V with respect to φ* , denoted by $G = H \rtimes_{\varphi} V$, is the Cartesian product $H \times V$ endowed with the group operation $*: G \times G \rightarrow G$ given by

$$(h_1, v_1) * (h_2, v_2) = (h_1 \cdot \varphi_{v_1}(h_2), v_1 v_2).$$

With this group operation, then both H and V can be viewed as subgroups of G and $H \triangleleft G$ is identified to a normal subgroup of G . This construction comes to generalize the notion of direct product of groups, where the operation on the cartesian product $H \times V$ would be the product operation $(h_1, v_1) * (h_2, v_2) = (h_1 h_2, v_1 v_2)$.

Even on the particular case of $H = \mathbb{R}^2$ and $V = \mathbb{R}$ being two abelian groups, it is possible to obtain a great variety of groups via the semidirect product of $\mathbb{R}^2$ and $\mathbb{R}$, depending uniquely on the choice of the (now 1-parameter) family of automorphisms of $\mathbb{R}^2$. Precisely, with the exceptions of $SU(2)$ (which is not diffeomorphic to $\mathbb{R}^3$) and $\widetilde{PSL}(2, \mathbb{R})$ (which has no normal subgroup of dimension 2), it is possible to construct all three dimensional simply connected Lie groups using the following setting: fix a matrix $A \in M_2(\mathbb{R})$,

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \tag{3}$$

and consider φ the 1 parameter subgroup of automorphisms of $\mathbb{R}^2$ generated by the exponential map

$$\begin{aligned}\varphi: \quad \mathbb{R} &\rightarrow \text{Aut}(\mathbb{R}^2) \\ z &\mapsto e^{Az}: \mathbb{R}^2 \rightarrow \mathbb{R}^2,\end{aligned}$$

then $\mathbb{R}^2 \rtimes_A \mathbb{R} = \mathbb{R}^2 \rtimes_{\varphi} \mathbb{R}$ is the semidirect product between $\mathbb{R}^2$ and $\mathbb{R}$ with respect to φ , i.e., the set $\mathbb{R}^3 = \mathbb{R}^2 \times \mathbb{R}$ endowed with the group operation $*$ defined via

$$(x_1, y_1, z_1) * (x_2, y_2, z_2) = \left(\begin{pmatrix} x_1 \\ y_1 \end{pmatrix} + e^{Az_1} \begin{pmatrix} x_2 \\ y_2 \end{pmatrix}, z_1 + z_2 \right). \quad (4)$$

Using the notation of [MP], denote the exponential map e^{Az} by

$$e^{Az} = \begin{pmatrix} a_{11}(z) & a_{12}(z) \\ a_{21}(z) & a_{22}(z) \end{pmatrix}, \quad (5)$$

and observe that the vector fields defined by

$$E_1(x, y, z) = a_{11}(z)\partial_x + a_{21}(z)\partial_y, \quad E_2(x, y, z) = a_{12}(z)\partial_x + a_{22}(z)\partial_y, \quad E_3 = \partial_z \quad (6)$$

are left invariant and extend the canonical basis $\{\partial_x(0), \partial_y(0), \partial_z(0)\}$ at the origin of $\mathbb{R}^3$. Moreover, if we let

$$F_1 = \partial_x, \quad F_2 = \partial_y, \quad F_3(x, y, z) = (ax + by)\partial_x + (cx + dy)\partial_y + \partial_z, \quad (7)$$

it follows that each F_i is a right invariant vector field of $\mathbb{R}^2 \rtimes_A \mathbb{R}$, so they are Killing fields with respect to any left invariant metric of $\mathbb{R}^2 \rtimes_A \mathbb{R}$.

The metric to be considered on $\mathbb{R}^2 \rtimes_A \mathbb{R}$ is the *canonical left invariant metric*, that is the one given by stating that $\{E_1, E_2, E_3\}$ are unitary and orthogonal to each other everywhere. In particular, as it holds

$$\begin{aligned}\partial_x(x, y, z) &= a_{11}(-z)E_1 + a_{21}(-z)E_2 \\ \partial_y(x, y, z) &= a_{12}(-z)E_1 + a_{22}(-z)E_2,\end{aligned}$$

we can express the metric of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ in coordinates as

$$\begin{aligned}ds^2 &= [a_{11}(-z)^2 + a_{21}(-z)^2] dx^2 + [a_{12}(-z)^2 + a_{22}(-z)^2] dy^2 + dz^2 \\ &\quad + [a_{11}(-z)a_{12}(-z) + a_{21}(-z)a_{22}(-z)] (dx \otimes dy + dy \otimes dx).\end{aligned}$$

Note that, as $e^{-Az} = (e^{Az})^{-1}$ and $\det(e^{Az}) = e^{z\text{trace}(A)}$, we have

$$\begin{pmatrix} a_{11}(-z) & a_{12}(-z) \\ a_{21}(-z) & a_{22}(-z) \end{pmatrix} = e^{-z\text{trace}(A)} \begin{pmatrix} a_{22}(z) & -a_{12}(z) \\ -a_{21}(z) & a_{11}(z) \end{pmatrix},$$

and we can introduce the notation

$$\begin{aligned} Q_{11}(z) &= \langle \partial_x, \partial_x \rangle = e^{-2z\text{trace}(A)} [a_{21}(z)^2 + a_{22}(z)^2] \\ Q_{22}(z) &= \langle \partial_y, \partial_y \rangle = e^{-2z\text{trace}(A)} [a_{11}(z)^2 + a_{12}(z)^2] \\ Q_{12}(z) &= \langle \partial_x, \partial_y \rangle = -e^{-2z\text{trace}(A)} [a_{11}(z)a_{21}(z) + a_{12}(z)a_{22}(z)] \end{aligned} \quad (8)$$

to obtain that the metric ds^2 is expressed by

$$ds^2 = Q_{11}(z)dx^2 + Q_{22}(z)dy^2 + dz^2 + Q_{12}(z)(dx \otimes dy + dy \otimes dx). \quad (9)$$

If $A, B \in M_2(\mathbb{R})$ are two congruent matrices, on the sense that there is some orthogonal matrix $P \in O(2)$ such that $B = PAP^{-1}$, then the groups $\mathbb{R}^2 \rtimes_A \mathbb{R}$ and $\mathbb{R}^2 \rtimes_B \mathbb{R}$, endowed with their respective canonical left invariant metrics are *isomorphic* and *isometric*, and the map that makes such identification is a simple rotation on horizontal planes induced by P ,

$$\begin{aligned} \psi: \quad \mathbb{R}^2 \rtimes_A \mathbb{R} &\rightarrow \mathbb{R}^2 \rtimes_B \mathbb{R} \\ (x, y, z) &\mapsto (P(x, y), z). \end{aligned} \quad (10)$$

The Lie brackets of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ are given by


$$[E_1, E_2] = 0, \quad [E_3, E_1] = aE_1 + cE_2, \quad [E_3, E_2] = bE_1 + dE_2, \quad (11)$$

so Levi-Civita equation implies that the Riemannian connection of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ satisfies

$$\begin{array}{l|l|l} \nabla_{E_1} E_1 & = aE_3 & \left| \begin{array}{l} \nabla_{E_1} E_2 = \frac{b+c}{2} E_3 \\ \nabla_{E_2} E_2 = dE_3 \end{array} \right| \begin{array}{l} \nabla_{E_1} E_3 = -aE_1 - \frac{b+c}{2} E_2 \\ \nabla_{E_2} E_3 = -\frac{b+c}{2} E_1 - dE_2 \end{array} \\ \nabla_{E_2} E_1 & = \frac{b+c}{2} E_3 & \\ \nabla_{E_3} E_1 & = \frac{c-b}{2} E_2 & \left| \begin{array}{l} \nabla_{E_3} E_2 = \frac{b-c}{2} E_1 \\ \nabla_{E_3} E_3 = 0. \end{array} \right. \end{array}$$

We notice two important properties of planes in $\mathbb{R}^2 \rtimes_A \mathbb{R}$: first, we observe that the metric ds^2 is invariant by rotations of angle π around the vertical lines $\{(x_0, y_0, z) \mid z \in \mathbb{R}\}$, hence vertical planes are minimal surfaces of $\mathbb{R}^2 \rtimes_A \mathbb{R}$. Moreover, horizontal planes $\{z = c\}$ have E_3 as an unitary normal vector field, so they have constant mean curvature (with respect to the upward orientation) given by $H = \text{trace}(A)/2$. In particular, horizontal planes of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ are going to be minimal if and only if $\text{trace}(A) = 0$.

However, the difference between the cases $\text{trace}(A) = 0$ and $\text{trace}(A) \neq 0$ go further than horizontal planes being minimal: concerning the classification of simply connected Lie groups of dimension 3, we notice that Meeks and Pérez, [MP] proved that any *non-unimodular*¹ Lie group of dimension 3



(a) A foliation of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ by vertical (minimal) planes (b) The foliation of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ by horizontal (CMC) planes

Figure 1: On semidirect products $\mathbb{R}^2 \rtimes_A \mathbb{R}$, every *vertical* plane is a minimal surface. Horizontal planes are flat, have constant mean curvature $H = \text{trace}(A)/2$ and the subgroup $\mathbb{H} = \mathbb{R}^2 \rtimes_A \{0\}$ (highlighted in the above right picture) is normal in $\mathbb{R}^2 \rtimes_A \mathbb{R}$.

is isomorphic and isometric to a semidirect product $\mathbb{R}^2 \rtimes_A \mathbb{R}$, endowed with its left invariant metric, where $A \in M_2(\mathbb{R})$ is such that $\text{trace}(A) \neq 0$ (Lemma 2.11, [MP]). Moreover, they also prove that, with the exceptions of $SU(2)$ and $\widehat{PSL}(2, \mathbb{R})$, all other unimodular metric Lie groups are isomorphic and isometric to a semidirect product $\mathbb{R}^2 \rtimes_A \mathbb{R}$, with $\text{trace}(A) = 0$ (Section 2.6 and Theorem 2.15, [MP]). Herein, we refer to the cases $\text{trace}(A) = 0$ or $\text{trace}(A) \neq 0$ respectively as the *unimodular* and *non-unimodular* case.

3. Mean curvature equation and height estimates

In this section, we consider a smooth open domain $\Omega \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ and a function $u: \Omega \rightarrow \mathbb{R}$. The π -graph of u is

$$\Sigma = \text{graph}(u) = \{(x, y, u(x, y)) \in \mathbb{R}^2 \rtimes_A \mathbb{R} \mid (x, y, 0) \in \Omega\}.$$

When oriented with respect to the upwards direction, the mean curvature of Σ is

$$H = \frac{e^{2\text{trace}(A)}}{2W^3} \left[u_{xx} (Q_{22}(u) + u_y^2) + u_{yy} (Q_{11}(u) + u_x^2) - 2u_{xy} (Q_{12}(u) + u_x u_y) + G_1(u)u_x^2 + G_2(u)u_y^2 + G_3(u)u_x u_y + (a + d)e^{-2\text{trace}(A)} \right], \quad (12)$$

¹A group G is said to be unimodular if $\det(\text{Ad}_g) = 1$ for all $g \in G$

where $Q_{ij}: \mathbb{R} \rightarrow \mathbb{R}$ are the coefficients of the metric of $\mathbb{R}^2 \rtimes_A \mathbb{R}$, defined in (8), $G_i: \mathbb{R} \rightarrow \mathbb{R}$ are the functions given by

$$\begin{aligned} G_1(z) &= e^{-2z\text{trace}(A)} \left((2a+d)a_{11}(z)^2 + (a+2d)a_{12}(z)^2 + (b+c)a_{11}(z)a_{12}(z) \right) \\ G_2(z) &= e^{-2z\text{trace}(A)} \left((2a+d)a_{21}(z)^2 + (a+2d)a_{22}(z)^2 + (b+c)a_{21}(z)a_{22}(z) \right) \\ G_3(z) &= e^{-2z\text{trace}(A)} \left((4a+2d)a_{11}(z)a_{21}(z) + (2a+4d)a_{12}(z)a_{22}(z) \right. \\ &\quad \left. + (b+c)(a_{11}(z)a_{22}(z) + a_{12}(z)a_{21}(z)) \right), \end{aligned} \quad (13)$$

and W is

$$\begin{aligned} W(z, p) &= \sqrt{1 + (a_{11}(z)p_1 + a_{21}(z)p_2)^2 + (a_{12}(z)p_1 + a_{22}(z)p_2)^2} \\ &= \sqrt{1 + e^{2z\text{trace}(A)} (Q_{22}(z)p_1^2 - 2Q_{12}(z)p_1p_2 + Q_{11}(z)p_2^2)}. \end{aligned}$$

Following the above notation, we define the *mean curvature operator*

$$\begin{aligned} Q(u) &= u_{xx} (Q_{22}(u) + u_y^2) + u_{yy} (Q_{11}(u) + u_x^2) + 2u_{xy} (Q_{12}(u) - u_x u_y) \\ &\quad + G_1(u)u_x^2 + G_2(u)u_y^2 + G_3(u)u_x u_y + (a+d)e^{-2u\text{trace}(A)}, \end{aligned} \quad (14)$$

so $\text{graph}(u)$ is a minimal surface of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ if and only if u satisfies $Q(u) = 0$ in Ω .

Note that Q is a quasilinear elliptic operator, as the matrix

$$\mathcal{Q}(z, p) = \begin{pmatrix} Q_{22}(z) + p_2^2 & Q_{12}(z) - p_1 p_2 \\ Q_{12}(z) - p_1 p_2 & Q_{11}(z) + p_1^2 \end{pmatrix} \quad (15)$$

is positive definite for every $z \in \mathbb{R}$ and $p = (p_1, p_2) \in \mathbb{R}^2$, as it is easy to see using the relation

$$Q_{11}(z)Q_{22}(z) - Q_{12}(z)^2 = e^{-2z\text{trace}(A)}.$$

In the papers of Meeks, Mira, Pérez and Ros, [MMPR, MMPR2, MMPR3], some work has been done in order to understand constant mean curvature π -graphs: the fact that $\mathbb{R}^2 \rtimes_A \mathbb{R}$ admits a foliation by parallel horizontal planes of constant mean curvature $H = \text{trace}(A)/2$ determines much of the structure of those graphs. For instance, using this property and the mean curvature comparison principle, they are able to prove

LEMMA 3.1 (Assertion 15.5, [MMPR3]). *Let $D \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ be a convex compact disk and let $C = \partial D$ be its boundary. Consider $\pi(x, y, z) = (x, y, 0)$ the vertical projection. If $\Gamma \subseteq \pi^{-1}(C)$ is a closed simple curve such that the*

projection $\pi: \Gamma \rightarrow C$ monotonically parametrizes² C and $h: \Gamma \rightarrow \mathbb{R}$ is the height function, let $c_0 = \inf_{\Gamma} h$ and $c_1 = \sup_{\Gamma} h$. If Σ is a compact minimal surface with $\partial\Sigma = \Gamma$, it follows:

1. If $\text{trace}(A) \geq 0$, then $\Sigma \subseteq \pi^{-1}(D) \cap \{z \geq c_0\}$;
2. If $\text{trace}(A) \leq 0$, then $\Sigma \subseteq \pi^{-1}(D) \cap \{z \leq c_1\}$.

In the particular case of graphs, Lemma 3.1 implies that a minimal graph over some smooth domain $\Omega \subseteq \mathbb{R}^2 \rtimes_A \{0\}$, compact and convex, satisfies the maximum principle if $\text{trace}(A) \leq 0$ and satisfies the minimum principle if $\text{trace}(A) \geq 0$, satisfying both only in the unimodular case. However, when $\text{trace}(A) > 0$ no uniform upper bound is obtained, neither a lower bound when $\text{trace}(A) < 0$. This motivates the search for height estimates for minimal graphs, which is next result. Perhaps, the proof of Theorem 3.2 is as interesting as the result itself, as it gives some understanding on the behaviour of the operator Q , on the many possible settings for the matrix A . Such properties will be used on the proof of Theorem 3.5, and also in Section 4 to obtain the existence of minimal Killing graphs that converge to the Scherk-like fundamental piece of Theorem 4.1.

THEOREM 3.2. *Let $A \in M_2(\mathbb{R})$ and let $\mathbb{R}^2 \rtimes_A \mathbb{R}$ be a semidirect product endowed with its canonical left invariant metric. Let $\Omega \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ be a bounded, convex domain and let $\alpha \in \mathbb{R}$ be any given constant. Then, there exists a constant $C(\alpha) = C(\text{diam}(\Omega), \alpha)$ such that for every u satisfying $Q(u) \geq 0$ and $\sup_{\partial\Omega} u \leq \alpha$, it holds that*

$$\sup_{\Omega} u \leq \alpha + C(\alpha). \quad (16)$$

In particular, there is a constant C depending on $\text{diam}(\Omega)$ and on $\sup_{\partial\Omega} u$ such that every $u: \Omega \rightarrow \mathbb{R}$ whose graph has nonnegative mean curvature function with respect to the upwards orientation satisfies

$$\sup_{\Omega} u \leq \sup_{\partial\Omega} u + C \left(\sup_{\partial\Omega} u \right). \quad (17)$$

The proof of Theorem 3.2 uses techniques from quasilinear elliptic partial differential equations, mainly the comparison principle. For instance, Theorem 10.1 of [GT] gives us that if R is a quasilinear elliptic operator of the form

$$R(w) = \sum_{i,j=1}^2 a_{ij}(x, \text{grad}(w)) w_{x_i x_j} + b(x, w, \text{grad}(w)), \quad (18)$$

²This means that $\pi(\Gamma) \subset \partial\Omega$ and $\pi^{-1}(\{p\}) \cap \Gamma$ is either a single point or a compact interval for every $p \in \partial\Omega$

for C^2 functions $w: \Omega \rightarrow \mathbb{R}$, where a_{ij} and b are smooth functions and b is such that for each $x \in \Omega$ and $p \in \mathbb{R}^2$ the function $z \mapsto b(x, z, p)$ is nonincreasing, then, given $u, v: \Omega \rightarrow \mathbb{R}$ such that $R(u) \geq R(v)$ in Ω and $u \leq v$ in $\partial\Omega$, then $u \leq v$ in Ω .

However, the operator Q given on by (14) does not satisfy the hypothesis of such comparison principle (or of its generalizations), as the coefficients of the second order terms of $Q(u)$ depend on u . This happens because translations $(x, y, z) \mapsto (x, y, z + t)$ are not isometries of $\mathbb{R}^2 \rtimes_A \mathbb{R}$. Hence, we are not able to prove uniqueness of solutions to the minimal graph equation and we also need to use an indirect approach to find the height estimates of Theorem 3.2.

In order to prove Theorem 3.2, we define a quasilinear operator R related to Q , for which it holds the comparison principle. Then, we find an *ad hoc* positive function $v: \Omega \rightarrow \mathbb{R}$, whose construction will depend only on Ω and on α such that $R(v + \alpha) \leq R(u)$. Then, as $u \leq \alpha \leq v + \alpha$ along $\partial\Omega$, it will follow that $u \leq v + \alpha$ in Ω , and we can let $C(\alpha, \Omega)$ be given by $C = \sup_{\Omega} v$.

PROOF (Proof of Theorem 3.2). First, we notice that when $\text{trace}(A) \leq 0$, the result is trivial with $C = 0$ and without the need for an α , by Lemma 3.1. Thus we will suppose that $\text{trace}(A) > 0$ and focus on the non-unimodular case. Without loss of generality, after a homothety of the metric we may assume that $\text{trace}(A) = 2$ and that A is written as

$$A = \begin{pmatrix} 1+a & b \\ c & 1-a \end{pmatrix}, \quad (19)$$

for some $a, b, c \in \mathbb{R}$. We divide the proof into two cases, starting when A is not a diagonal matrix.

CASE 1. Suppose that A is not a diagonal matrix.

We begin by proving the following key claim, which will also be used in Section 4:

CLAIM 1. Let the functions Q_{ij} be the ones defined by (8) with respect to the matrix A of (19), where either $b \neq 0$ or $c \neq 0$. Then, there is some $\lambda > 0$ such that at least one of the following hold, for every $z \in \mathbb{R}$:

- i. $Q_{22}(z)e^{2z} > \lambda$;
- ii. $Q_{11}(z)e^{2z} > \lambda$.

Moreover, if $a^2 + bc \leq 0$, both i. and ii. hold, and if $a^2 + bc > 0$, then $b \neq 0$ is equivalent to i. and $c \neq 0$ is equivalent to ii.

Proof of Claim 1. We prove Claim 1 in each of three (family of) possibilities to the exponential of A . First, write $A = I + A_0$, where I is the identity matrix and A_0 is the traceless part of A ,

$$A_0 = \begin{pmatrix} a & b \\ c & -a \end{pmatrix}.$$

As I and A_0 commute, we obtain that $e^{Az} = e^{Iz+A_0z} = e^{Iz}e^{A_0z}$, thus

$$e^{Az} = e^z \begin{pmatrix} a_{11}^0(z) & a_{12}^0(z) \\ a_{21}^0(z) & a_{22}^0(z) \end{pmatrix},$$

where we denote by $a_{ij}^0(z)$ the coefficients of the exponential e^{A_0z} . Then $a_{ij}(z) = e^z a_{ij}^0(z)$, and it follows that

$$Q_{11}(z)e^{2z} = e^{-4z} [a_{21}(z)^2 + a_{22}(z)^2] e^{2z} = a_{21}^0(z)^2 + a_{22}^0(z)^2, \quad (20)$$

and, analogously,

$$Q_{22}(z)e^{2z} = a_{11}^0(z)^2 + a_{12}^0(z)^2. \quad (21)$$

Note that the characteristic equation of A_0 is $0 = \det(A_0 - tI) = t^2 - (a^2 + bc)$, so if let $d = \sqrt{|a^2 + bc|}$, the exponential of A_0 is³

$$e^{A_0z} = \begin{pmatrix} \cos(dz) + \frac{a}{d} \sin(dz) & \frac{b}{d} \sin(dz) \\ \frac{c}{d} \sin(dz) & \cos(dz) - \frac{a}{d} \sin(dz) \end{pmatrix}, \text{ when } a^2 + bc < 0, \quad (22)$$

$$e^{A_0z} = \begin{pmatrix} 1 + az & bz \\ cz & 1 - az \end{pmatrix}, \text{ when } a^2 + bc = 0, \quad (23)$$

$$e^{A_0z} = \begin{pmatrix} \cosh(dz) + \frac{a}{d} \sinh(dz) & \frac{b}{d} \sinh(dz) \\ \frac{c}{d} \sinh(dz) & \cosh(dz) - \frac{a}{d} \sinh(dz) \end{pmatrix}, \text{ when } a^2 + bc > 0. \quad (24)$$

Let $f(z) = a_{11}^0(z)^2 + a_{12}^0(z)^2$ and $g(z) = a_{21}^0(z)^2 + a_{22}^0(z)^2$. We will prove that there is some $\lambda > 0$ such that either $f(z) > \lambda$ or $g(z) > \lambda$, and this proves the claim, by (20) and (21).

Note that both f and g are always positive, as the existence of some $z_0 \in \mathbb{R}$ such that $f(z_0) = 0$ or $g(z_0) = 0$ would imply that $\det(e^{A_0z_0}) = 0$, an absurdity. Hence, we just need to check the asymptotic behaviour of f and g .

³We remark that the constant $a^2 + bc$ is linked with the Milnor D -invariant of $\mathbb{R}^2 \rtimes_A \mathbb{R}$, which is defined by $D = \det(A) = 1 - (a^2 + bc)$. So each case $a^2 + bc > 0$, $a^2 + bc = 0$ and $a^2 + bc < 0$ is in correspondence with $D < 1$, $D = 1$ and $D > 1$, respectively.

If $a^2 + bc < 0$, the existence of λ as claimed follows directly from the fact that both f and g are periodic and positive, by (22). If $a^2 + bc = 0$, then we have that f and g are

$$\begin{aligned} f(z) &= (1 + az)^2 + (bz)^2 = (a^2 + b^2)z^2 + 2az + 1 \\ g(z) &= (1 - az)^2 + (cz)^2 = (a^2 + c^2)z^2 - 2az + 1, \end{aligned}$$

both strictly positive at infinity for any choice of a, b, c , so we also have the existence of λ in this case. Finally, if $a^2 + bc > 0$, f and g are

$$\begin{aligned} f(z) &= \left(\cosh(dz) + \frac{a}{d} \sinh(dz) \right)^2 + \left(\frac{b}{d} \sinh(dz) \right)^2 \\ g(z) &= \left(\cosh(dz) - \frac{a}{d} \sinh(dz) \right)^2 + \left(\frac{c}{d} \sinh(dz) \right)^2. \end{aligned}$$

If $i.$ was not true, either $\lim_{z \rightarrow -\infty} f(z) = 0$ or $\lim_{z \rightarrow +\infty} f(z) = 0$, hence $b = 0$. Also, if $\lim_{z \rightarrow -\infty} g(z) = 0$ or $\lim_{z \rightarrow +\infty} g(z) = 0$, we would have $c = 0$. This shows that if $b \neq 0$, then $i.$ holds, and if $c \neq 0$ then $ii.$ holds. As A is not a diagonal matrix, at least one between $i.$ and $ii.$ is true, finishing the proof of the claim. $\diamond$

To proceed with the proof of the first case of Theorem 3.2, we prove the existence of $\Lambda > 0$ such that $G_1(z) \leq \Lambda Q_{22}(z)$ and $G_2(z) \leq \Lambda Q_{11}(z)$. By definition,

$$\begin{aligned} \frac{G_1(z)}{Q_{22}(z)} &= \frac{e^{-4z} [(3+a)a_{11}(z)^2 + (3-a)a_{12}(z)^2 + (b+c)a_{11}(z)a_{12}(z)]}{e^{-4z} [a_{11}(z)^2 + a_{12}(z)^2]} \\ &= 3 + a \frac{a_{11}(z)^2 - a_{12}(z)^2}{a_{11}(z)^2 + a_{12}(z)^2} + (b+c) \frac{a_{11}(z)a_{12}(z)}{a_{11}(z)^2 + a_{12}(z)^2} \\ &\leq 3 + |a| + \frac{|b+c|}{2} = \Lambda, \end{aligned} \tag{25}$$

and, mutatis mutandis, the same estimate holds for the quotient $G_2(z)/Q_{11}(z)$.

Next, using the existence of λ and Λ as before we prove the first case of the theorem. Fix any constant $\alpha \in \mathbb{R}$ and let u be any function that satisfies $Q(u) \geq 0$ and $\sup_{\partial\Omega} u \leq \alpha$.

First, assume that $i.$ holds and let R be the quasilinear elliptic operator defined as

$$\begin{aligned} R(w) &= w_{xx} \left(\frac{Q_{22}(u) + w_y^2}{Q_{22}(u)} \right) + w_{yy} \left(\frac{Q_{11}(u) + w_x^2}{Q_{22}(u)} \right) + 2w_{xy} \left(\frac{Q_{12}(u) - w_x w_y}{Q_{22}(u)} \right) \\ &\quad + \frac{G_1(u)}{Q_{22}(u)} w_x^2 + \frac{G_2(u)}{Q_{22}(u)} w_y^2 + \frac{G_3(u)}{Q_{22}(u)} w_x w_y + 2 \frac{e^{-2u}}{Q_{22}(u)} e^{-2w}. \end{aligned} \tag{26}$$

Note that R is defined in order to have two features. First, when $w = u$, we have $R(u) = Q(u)/Q_{22}(u) \geq 0$. Second, using the notation of (18), we have that the coefficients a_{ij} of R do not depend on w , only on the space variable and on the derivatives of w . Also, the function $z \mapsto b(x, z, p)$ is nonincreasing for every x and p fixed, thus R satisfies the hypothesis of the comparison principle (although, as noticed before, Q does not).

In order to finish the proof of Case 1 (when $i.$ holds), we will build a nonnegative function $v: \Omega \rightarrow \mathbb{R}$ that will depend uniquely on Ω and on α such that $R(v + \alpha) \leq 0 \leq R(u)$. As $u \leq \alpha \leq \alpha + v$ on $\partial\Omega$, it will follow from the comparison principle that $u \leq v + \alpha$ in Ω , and this will finish the proof.

As Ω is a bounded domain, after a horizontal translation (which is an isometry of the ambient space) we may assume without loss of generality that it is contained in a strip

$$\Omega \subseteq \{(x, y, 0) \in \mathbb{R}^2 \rtimes_A \mathbb{R} \mid 1 < x < M\},$$

for some $M > 1$. Let $v(x, y) = \ln(lx)/L$, where $l, L > 0$ are constants yet to be defined. By the definition of R and v , and by the existence of λ and Λ as before, we have that

$$\begin{aligned} R(v + \alpha) &= v_{xx} + \frac{G_1(u)}{Q_{22}(u)} v_x^2 + 2 \frac{e^{-2u}}{Q_{22}(u)} e^{-2(v+\alpha)} \\ &< v_{xx} + \Lambda v_x^2 + \frac{2}{\lambda} e^{-2(v+\alpha)}. \end{aligned}$$

Then, using that $e^v = (lx)^{\frac{1}{L}}$, $v_x = \frac{1}{Lx}$ and $v_{xx} = \frac{-1}{Lx^2}$, we obtain

$$\begin{aligned} R(v + \alpha) &< -\frac{1}{Lx^2} + \Lambda \frac{1}{L^2 x^2} + \frac{2}{\lambda e^{2\alpha}} (lx)^{-2/L} \\ &= \frac{1}{Lx^2} \left[-1 + \frac{\Lambda}{L} + \frac{2L}{\lambda e^{2\alpha} l^{2/L}} x^{(2L-2)/L} \right]. \end{aligned} \tag{27}$$

Take $L = 1 + \Lambda$. As $1 < x < M$, it follows that

$$R(v + \alpha) < \frac{1}{(1 + \Lambda)x^2} \left[-\frac{1}{1 + \Lambda} + 2 \frac{1 + \Lambda}{\lambda e^{2\alpha} l^{\frac{2}{1+\Lambda}}} M^{\frac{2\Lambda}{1+\Lambda}} \right], \tag{28}$$

and we can choose l big enough (in particular we may assume $l \geq 1$, so that $v > 0$ in Ω) such that

$$-\frac{1}{1 + \Lambda} + 2 \frac{1 + \Lambda}{\lambda e^{2\alpha} l^{\frac{2}{1+\Lambda}}} M^{\frac{2\Lambda}{1+\Lambda}} < 0, \tag{29}$$

so $R(v + \alpha) < 0$. We remark that the choice of l and L as above depends uniquely on λ, Λ, α and M , so it does not depend on u .

As R satisfies the hypothesis of the Comparison Principle and $v + \alpha \geq u$ on $\partial\Omega$, it follows that $\sup_{\Omega} u \leq \sup_{\Omega} v + \alpha$. Finally, we set $C = \sup_{\Omega} v$, and the theorem follows when A is not diagonal and *i.* holds.

Still in Case 1 of A not being a diagonal matrix, if *i.* was not true, then $b = 0$, and $c \neq 0$, then *ii.* holds. In this case, let

$$\begin{aligned} R(w) = & w_{xx} \left(\frac{Q_{22}(u) + w_y^2}{Q_{11}(u)} \right) + w_{yy} \left(\frac{Q_{11}(u) + w_x^2}{Q_{11}(u)} \right) + 2w_{xy} \left(\frac{Q_{12}(u) - w_x w_y}{Q_{11}(u)} \right) \\ & + \frac{G_1(u)}{Q_{11}(u)} w_x^2 + \frac{G_2(u)}{Q_{11}(u)} w_y^2 + \frac{G_3(u)}{Q_{11}(u)} w_x w_y + 2 \frac{e^{-2u}}{Q_{11}(u)} e^{-2w}. \end{aligned} \quad (30)$$

From here, just proceed analogously as before, however using $v(x, y) = \ln(l y)/L$, and making the appropriate choices to l and L to finish the proof of Case 1.

CASE 2. Assume that A is a diagonal matrix

$$A = \begin{pmatrix} 1+a & 0 \\ 0 & 1-a \end{pmatrix}.$$

In this case, $a_{11}(z) = e^{(1+a)z}$, $a_{22}(z) = e^{(1-a)z}$ and $a_{12}(z) = a_{21}(z) = 0$. It follows that the operator Q is

$$\begin{aligned} Q(u) = & u_{xx} \left(e^{-2(1-a)u} + u_y^2 \right) + u_{yy} \left(e^{-2(1+a)u} + u_x^2 \right) - 2u_{xy} (u_x u_y) \\ & + (3+a)e^{-2(1-a)u} u_x^2 + (3-a)e^{-2(1+a)u} u_y^2 + 2e^{-4u}. \end{aligned}$$

If $a \geq 0$ we define R as the operator

$$\begin{aligned} R(w) = & w_{xx} \left(1 + e^{2(1-a)u} w_y^2 \right) + w_{yy} \left(e^{-4au} + e^{2(1-a)u} w_x^2 \right) - 2w_{xy} \left(e^{2(1-a)u} w_x w_y \right) \\ & + (3+a)w_x^2 + (3-a)e^{-4au} w_y^2 + 2e^{-2(1+a)u}, \end{aligned} \quad (31)$$

and if $a < 0$, R will be defined as

$$\begin{aligned} R(w) = & w_{xx} \left(e^{4au} + e^{2(1+a)u} w_y^2 \right) + w_{yy} \left(1 + e^{2(1+a)u} w_x^2 \right) - 2w_{xy} \left(e^{2(1+a)u} w_x w_y \right) \\ & + (3+a)e^{4au} w_x^2 + (3-a)w_y^2 + 2e^{-2(1-a)u}. \end{aligned} \quad (32)$$

Now, we just set v to be again $v(x, y) = \ln(l x)/L$ when $a \geq 0$ and $v(x, y) = \ln(l y)/L$ when $a < 0$ and the proof will follow as in the previous case, using $\Lambda = 3 + |a|$ and $\lambda = 1$. $\square$

Next theorem is to prove that the dependence on α on the constant C of Theorem 3.2 cannot be removed. Precisely, we prove

THEOREM 3.3. *Let A be a matrix as in (19) and let $X = \mathbb{R}^2 \rtimes_A \mathbb{R}$ be a non-unimodular semidirect product endowed with its canonical left invariant metric. Let $\Omega \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ be a bounded, convex domain. Then, for every constant $C > 0$ there exists some function $u: \Omega \rightarrow \mathbb{R}$ satisfying $Q(u) = 0$ and also*

$$\sup_{\Omega} u > \sup_{\partial\Omega} u + C. \quad (33)$$

The proof of Theorem 3.3 above is by contradiction and consists on using the vertical translation that rises from the group structure to translate a family of solutions tending to $-\infty$, all to height 0. We prove that if Theorem 3.3 was false, such family would be uniformly bounded, and this would generate a contradiction with the following theorem, due to Meeks, Mira, Pérez and Ros [MMPR3]

THEOREM 3.4 (Theorem 15.4, [MMPR3]). *Let X be a non-unimodular metric Lie group which is isomorphic and isometric to a semidirect product $\mathbb{R}^2 \rtimes_A \mathbb{R}$, $A \in M_2(\mathbb{R})$. Suppose that $\Gamma(n) \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ is a sequence of C^2 simple closed convex curves with $e = (0, 0, 0) \in \Gamma(n)$ such that the geodesic curvatures of $\Gamma(n)$ converge uniformly to 0 and the curves $\Gamma(n)$ converge on compact subsets to a line L with $e \in L$ as $n \rightarrow \infty$. Then, for any sequence $M(n)$ of compact branched minimal disks with $\partial M(n) = \Gamma(n)$, the surfaces $M(n)$ converge C^2 on compact subsets as $n \rightarrow \infty$ to the vertical half plane $\pi^{-1}(L) \cap [\mathbb{R}^2 \rtimes_A [0, \infty)]$.*

PROOF (Proof of Theorem 3.3). We begin by proving the following claim:

CLAIM 2. Let $\mathbb{S}^1 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$ be the unit circle centred on the origin of $\mathbb{R}^2$. Let $A \in M_2(\mathbb{R})$ be a matrix with $\text{trace}(A) = 2$, as in (19), and let e^{Az} be its exponential map. Then, there is a point $p \in \mathbb{S}^1$ and an increasing sequence $(z_n)_{n \in \mathbb{N}} \in (0, +\infty)$ such that the curves $\Gamma_n = e^{Az_n}(\mathbb{S}^1 - p)$, defined by the homothety by e^{Az_n} of the translated circle $\mathbb{S}^1 - p$, satisfies the hypothesis of Theorem 3.4 at the origin, i.e., the geodesic curvature of Γ_n at 0 is converging to zero and Γ_n is converging to a line L on compacts, with $0 \in L$.

Proof of Claim 2. Denote by A_0 the traceless part of A and observe that $e^{Az} = e^z e^{A_0 z}$. Then we have that $e^{Az} \mathbb{S}^1 = e^z (e^{A_0 z} \mathbb{S}^1)$ is a homothety by e^z of the curve $e^{A_0 z} \mathbb{S}^1$. Let $d = \sqrt{|a^2 + bc|}$ and divide the proof on the three aforementioned cases given on (22), (23) and (24).

If $a^2 + bc < 0$, we let $p \in \mathbb{S}^1$ be any point and define $z_n = \frac{2n\pi}{d}$. Then $e^{A_0 z_n} = \text{Id}$, so $e^{Az_n} \mathbb{S}^1$ is a circle of radius e^{2z_n} centred at the origin, and

$\Gamma_n = e^{Az_n}(\mathbb{S}^1 - p)$ is a circle through the origin with radius e^{2z_n} . As $z_n \rightarrow \infty$, Γ_n will converge to a line L through 0 and the claim is proved in this case.

If $a^2 + bc = 0$, then $e^{A_0 z}$ is given by (23) and $e^{A_0 z}\mathbb{S}^1$ is an ellipse and the homotheties of an ellipse by e^n admits a point where its geodesic curvature converges to zero and, after a translation, it converges to a line on compacts, proving the claim in this second case.

Finally, if $a^2 + bc > 0$, $e^{A_0 z}$ is given by (24). If $bc \neq 0$, then $d \neq |a|$, and if z is big enough we have that $\cosh(dz) \simeq e^{dz}/2$ and $\sinh(dz) \simeq e^{dz}/2$, so

$$e^{A_0 z} \simeq \frac{e^{dz}}{2d} \begin{pmatrix} d+a & b \\ c & d-a \end{pmatrix},$$

and $e^{Az}\mathbb{S}^1$ is asymptotic to a homothety of $e^{(d+2)z}$ of an ellipse, which has the desired properties. The last case to be treated is when $d^2 = a^2 + bc = a^2 > 0$, then

$$e^{A_0 z} = \begin{pmatrix} e^{dz} & \frac{b}{d} \sinh(dz) \\ \frac{c}{d} \sinh(dz) & e^{-dz} \end{pmatrix} \simeq \frac{e^{dz}}{d} \begin{pmatrix} d & b \\ c & de^{-2dz} \end{pmatrix},$$

and, for z large enough it follows that $e^{A_0 z}\mathbb{S}^1$ is asymptotic to a line segment, with multiplicity 2. Now, it depends on the two possible cases $0 < d \leq 1$ or $d > 1$ to understand what is the convergence of $e^{Az}\mathbb{S}^1$: if $d \leq 1$, then the homothety of e^z on $e^{A_0 z}$ will open the segment and make it asymptotic to an ellipse, which again admits a point p as claimed. If $d > 1$, then the action of e^z still makes $e^{Az}\mathbb{S}^1$ converge to a line on compacts, so the claim is proved. $\diamond$

Now, we prove Theorem 3.3 arguing by contradiction. Suppose that there is $C > 0$ such that, for every solution of $Q(u) = 0$ in Ω , it holds

$$\sup_{\Omega} u \leq \sup_{\partial\Omega} u + C. \quad (34)$$

In particular, the same estimate holds for any bounded, smooth domain contained in Ω . Let $r > 0$ be such that an euclidean ball B_r with radius r is contained in Ω and let $\mathbb{S}^1(r) = \partial B_r$ be the circle that bounds B_r and let $p \in \mathbb{S}^1(r)$ and $(z_n)_{n \in \mathbb{N}}$ be the ones given by Claim 2. Consider, for each $n \in \mathbb{N}$, the problem

$$\begin{cases} Q(u) = 0 & \text{in } B_r \\ u = -z_n & \text{on } \partial B_r. \end{cases} \quad (35)$$

The existence result due to Meeks, Mira, Pérez and Ros, Theorem 15.1 of [MMPR3], implies that (35) admits a solution $u_n: B_r \rightarrow \mathbb{R}$, and, from (34), it follows that, for every $n \in \mathbb{N}$, u_n satisfies

$$\sup_{B_r} u_n \leq \sup_{\partial\Omega} u_n + C = -z_n + C.$$

We translate the functions u_n vertically to height 0, using the left translation of the group $L_{(0,0,z_n)}$ to obtain a contradiction. If $\Sigma_n = \text{graph}(u_n)$, then

$$\begin{aligned} L_{(0,0,z_n)}\Sigma_n &= \left\{ L_{(0,0,z_n)}(x, y, u_n(x, y)) \mid (x, y) \in B_r \right\} \\ &= \left\{ \left(e^{Az_n} \begin{pmatrix} x \\ y \end{pmatrix}, u_n(x, y) + z_n \right), (x, y) \in B_r \right\} \\ &= \left\{ \left(\tilde{x}, \tilde{y}, u_n \left(e^{-Az_n} \begin{pmatrix} \tilde{x} \\ \tilde{y} \end{pmatrix} \right) + z_n \right), (\tilde{x}, \tilde{y}) \in e^{Az_n} B_r \right\}. \end{aligned}$$

Hence, if we let $v_n: e^{Az_n} B_r \rightarrow \mathbb{R}$ be the function defined by

$$v_n(x, y) = u_n \left(e^{-Az_n} \begin{pmatrix} x \\ y \end{pmatrix} \right) + z_n,$$

it follows that the graph of v_n is a left translate of the graph of u_n , in particular its graph $\tilde{\Sigma}_n = L_{(0,0,z_n)}\Sigma_n$ is a minimal surface of $\mathbb{R}^2 \rtimes_A \mathbb{R}$. Moreover, these graphs $\tilde{\Sigma}_n$ satisfy the hypothesis of Theorem 3.4, thus they should converge, in compact sets, to a vertical half plane. However, it holds that

$$\sup_{e^{Az_n} B_r} v_n = \sup_{B_r} u_n + z_n \leq C,$$

so the sequence v_n is uniformly bounded, generating a contradiction that proves Theorem 3.3. $\square$

Note that last proof shows more than the existence of a function u as on (33) for a fixed constant C . We actually proved that *any* sequence of functions with values along the boundary converging to $-\infty$ should have unbounded oscillation. In particular, using the notation of Theorem 3.2, it follows that, when $\alpha \rightarrow -\infty$, necessarily $C(\alpha) \rightarrow +\infty$. It is also possible to prove that $C(\alpha)$ may be chosen more carefully to satisfy $C(\alpha) \rightarrow 0$ when $\alpha \rightarrow +\infty$ (when $\text{trace}(A) > 0$). We make this analysis in the next result and in Corollary 3.6.

THEOREM 3.5. *Let $A \in M_2(\mathbb{R})$ and let $\mathbb{R}^2 \rtimes_A \mathbb{R}$ be a semidirect product endowed with its canonical left invariant metric. Let $\Omega \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ be some open, bounded, smooth domain, $k \in \mathbb{Z}$ be given and let u_k be a solution to the problem*

$$\begin{cases} Q(u) = 0 & \text{in } \Omega \\ u = k & \text{on } \partial\Omega. \end{cases} \quad (36)$$

Then, if $\text{osc}_\Omega(u) = \sup_\Omega(u) - \inf_\Omega(u)$ denotes the oscillation of a function u in Ω , we have

1. If $\text{trace}(A) = 0$, $u_k \equiv k$ is the constant function.

2. If $\text{trace}(A) > 0$, then $u_k > k$ in Ω . Moreover,

$$\lim_{k \rightarrow +\infty} \text{osc}_\Omega(u_k) = +\infty \text{ and } \lim_{k \rightarrow +\infty} \text{osc}_\Omega(u_k) = 0.$$

3. If $\text{trace}(A) < 0$, then $u_k < k$ in Ω . Moreover,

$$\lim_{k \rightarrow -\infty} \text{osc}_\Omega(u_k) = 0 \text{ and } \lim_{k \rightarrow +\infty} \text{osc}_\Omega(u_k) = +\infty.$$

PROOF. If $\text{trace}(A) = 0$, it is clear that $u_k \equiv k$ is the unique solution to (36), by Lemma 3.1, proving 1. Also, as the change $A \rightarrow -A$ gives rise to an isometry $(x, y, z) \in \mathbb{R}^2 \rtimes_A \mathbb{R} \mapsto (-x, -y, -z) \in \mathbb{R}^2 \rtimes_{-A} \mathbb{R}$, 3 follows from 2, so we can simply prove the case of $\text{trace}(A) > 0$, and, as previous, it is without loss of generality that we assume that $\text{trace}(A) = 2$, so A is written as in (19).

From Lemma 3.1, it follows that $u_k \geq k$ in Ω , and, if at an interior point $x \in \Omega$ the function u_k attains its minimum $u_k(x) = k$, then the mean curvature comparison principle, applied to $\Sigma_k = \text{graph}(u_k)$ and to the plane $\{z = k\}$ will imply that the mean curvature of Σ_k is at least as big as the one of the plane, which is $1 > 0$, a contradiction that proves that $u_k > k$ in Ω .

The second part of 2 follows like on the proof of Theorem 3.3: if the oscillation of u_k was not going to $+\infty$ when $k \rightarrow -\infty$ then we could translate all the minimal surfaces $\Sigma_k = \text{graph}(u_k)$ to height zero and obtain a contradiction with Theorem 3.4.

It remains to prove that the oscillation of u_k goes to zero when k approaches $+\infty$. In order to do so, it suffices to prove that the constant $C(\alpha)$ goes to zero when $\alpha \rightarrow \infty$.

Recall the proof of Theorem 3.2: $C = C(\alpha)$ was chosen depending on many parameters $l, L, \lambda, \Lambda, M, \alpha$. The constants λ and Λ depend only on the ambient space, as they come from estimates to the coefficients of the operator Q . The constant M depends uniquely on the diameter of Ω , so it is also fixed. In the proof of Theorem 3.2, the free parameters we could work with were l and L , depending on the previous ones and on the a priori constant α . Using an appropriate choice of l and L , we obtained the following expression to C

$$C = \frac{\ln(lM)}{L}.$$

The key steps to chose l and L were between equations (27), (28) and (29). However, these steps were done by thinking on the worst case, where the number α was a *negatively large* number, so we began by choosing L and then got to the definition of a l big enough, in order to compensate $e^{2\alpha}$,

which was though to be close to zero. Now, we are taking $\alpha_k = k$ to be *positively large*, so we follow a different approach.

Using the notation of the proof of Theorem 3.2, let $L = \Lambda + j$, where $j \in \mathbb{N}$ is yet to be chosen, and take $l = 1$, to obtain, similarly to (28), the inequality

$$R(v+k) < \frac{1}{(\Lambda+j)x^2} \left[-\frac{j}{\Lambda+j} + 2\frac{\Lambda+j}{\lambda e^{2k}} M^{\left(2-\frac{2}{\Lambda+j}\right)} \right]. \quad (37)$$

Then, we proceed as before, and try to find some $j \in \mathbb{N}$ such that the right hand side of (37) becomes negative. Such j exists if and only if it satisfies

$$\frac{(\Lambda+j)^2}{jM^{\frac{2}{\Lambda+j}}} < \frac{\lambda}{2M^2} e^{2k}. \quad (38)$$

There is $k_0 \in \mathbb{N}$ big enough such that for every $k \geq k_0$ it is possible to find $j \in \mathbb{N}$ satisfying (38). For $k \geq k_0$, denote by $j(k)$ the *largest* $j \in \mathbb{N}$ such that (38) hold (as the left hand side is unbounded with j this is well defined). By taking $L = \Lambda + j(k)$, we use (37) to obtain, as in Theorem 3.2, that exists a constant $C(k) = C(\Omega, k)$ given by

$$C(k) = \frac{\ln(M)}{\Lambda + j(k)}$$

such that every $u: \Omega \rightarrow \mathbb{R}$ such that

$$\begin{cases} Q(u) \geq 0 \text{ in } \Omega \\ u \leq k \text{ on } \partial\Omega \end{cases}$$

satisfies

$$\sup_{\Omega} u \leq k + \frac{\ln(M)}{\Lambda + j(k)}.$$

Note this is the same result as on Theorem 3.2 but for a different constant C , and only for $\alpha = k \geq k_0$. In particular, the functions u_k satisfy, for k large enough, that

$$\sup_{\Omega} u_k \leq k + \frac{\ln(M)}{\Lambda + j(k)},$$

hence

$$\text{osc}_{\Omega}(u_k) = \sup_{\Omega} u_k - k \leq \frac{\ln(M)}{\Lambda + j(k)}.$$

Finally, as the right hand side of (38) is unbounded with respect to k , it follows that $\lim_{k \rightarrow \infty} j(k) = +\infty$, so

$$\lim_{k \rightarrow +\infty} \frac{\ln(M)}{\Lambda + j(k)} = 0,$$

and it follows that the oscillation of u_k also tends to zero when $k \rightarrow +\infty$, finishing the proof of 2 and of the theorem. $\square$

This previous proof has as a consequence the next result.

COROLLARY 3.6. *Let $\mathbb{R}^2 \rtimes_A \mathbb{R}$ be a non-unimodular semidirect product with $\text{trace}(A) > 0$ and let $C(\alpha)$ be the constant given by Theorems 3.2 and by the proof of Theorem 3.5. Then*

$$\lim_{\alpha \rightarrow -\infty} C(\alpha) = +\infty, \quad \lim_{\alpha \rightarrow +\infty} C(\alpha) = 0.$$

In particular, if $u_L: \Omega \rightarrow \mathbb{R}$ is a function satisfying

$$\begin{cases} Q(u) \geq 0 \text{ on } \Omega, \\ \sup_{\partial\Omega} u = L \in \mathbb{R}. \end{cases}$$

Then

$$\lim_{L \rightarrow -\infty} \left(\sup_{\Omega} u_L - L \right) = +\infty, \quad \lim_{L \rightarrow +\infty} \left(\sup_{\Omega} u_L - L \right) = 0. \quad (39)$$

4. Scherk-like fundamental pieces

In this section, we use the tools developed in the study of the mean curvature operator, together with Killing graphs techniques to obtain an existence result of *Scherk-like fundamental pieces*, which are minimal π -graphs on $\mathbb{R}^2 \rtimes_A \mathbb{R}$ assuming the value 0 along a piecewise smooth curve $\gamma \subset \mathbb{R}^2 \rtimes_A \{0\}$ and having $\gamma \cup (\{p_1\} \times [0, \infty)) \cup (\{p_2\} \times [0, \infty))$ as boundary, where p_1 and p_2 are the endpoints of γ .

In the ambient space of an *unimodular* group $\mathbb{R}^2 \rtimes_A \mathbb{R}$, A. Menezes [Me] proved the existence of *complete* (without boundary) minimal surfaces, similar to the singly and to the doubly periodic Scherk minimal surfaces of $\mathbb{R}^3$. We would like to take a moment to give the main steps of the proof of Menezes to the existence of a doubly periodic example:

PROOF (Sketch of the proof of Theorem 2 of [Me]). Let $\Delta \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ be a triangle with vertexes

$$o = (0, 0, 0), \quad p_1 = (a, 0, 0), \quad p_2 = (0, a, 0),$$

for some $a > 0$. Let P_c be the polygon given by the union of segments

$$P_c = \overline{op_1} \cup \overline{p_1 p_1(c)} \cup \overline{p_1(c) p_2(c)} \cup \overline{p_2(c) p_2} \cup \overline{p_2 o}, \quad (40)$$

where $p_1(c) = (a, 0, c)$ and $p_2(c) = (0, a, c)$. Theorem 15.1 of [MMPR3] implies the existence of a minimal π -graph Σ_c with $\partial\Sigma_c = P_c$.

Then, one key property was observed: any Σ with such boundary is a *Killing graph* over the vertical domain $\Omega_c = \{(t, a - t, s) \mid 0 \leq t \leq a, 0 \leq s \leq c\}$ with respect to the horizontal Killing field $\partial_x + \partial_y$, thus Σ_c is the *unique* minimal surface with Γ_c as boundary.

This implies that Σ_c is stable and that the variation $c \mapsto \Sigma_c$ is continuous. By making $c \rightarrow \infty$, and using curvature estimates due to H. Rosenberg, R. Souam and E. Toubiana [RST] for stable surfaces in homogeneous manifolds, it is possible to show the convergence of Σ_c to some surface Σ_∞ , nowhere vertical and with boundary

$$\partial\Sigma_\infty = P_\infty = \overline{op_1} \cup (\{p_1\} \times [0, \infty)) \cup \overline{op_2} \cup (\{p_2\} \times [0, \infty)).$$

Finally, use the ambient isometries to rotate Σ_∞ along the two segments $\overline{op_1}$ and $\overline{op_2}$ to obtain a complete minimal π -graph on $\mathbb{R}^2 \rtimes_A \mathbb{R}$, which can be extended periodically by horizontal translations. $\square$

In this subject, our contribution is an extension of the above result to any semidirect product $\mathbb{R}^2 \rtimes_A \mathbb{R}$. Although in the general case our method does not produce examples without boundary, on the setting of unimodular groups our proof, which differs from the one of Menezes, reobtains the same result explained above. We state our result as follows.

THEOREM 4.1. *Let $\mathbb{R}^2 \rtimes_A \mathbb{R}$ be a semidirect product, where $A \in M_2(\mathbb{R})$ is any matrix with $\text{trace}(A) \geq 0$. Then, there is $L_0 = L_0(\text{trace}(A), \det(A)) > 0$ (and $L_0 = \infty$ when $\text{trace}(A) = 0$) such that if $p_1, p_2 \in \mathbb{R}^2 \rtimes_A \{0\}$ satisfy $d(p_1, p_2) \leq L_0$, then for any piecewise smooth curve $\gamma \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ with endpoints p_1, p_2 which is a convex graph over the segment $\alpha = \overline{p_1 p_2}$ and meets α on angles less than $\pi/2$, there exists a minimal surface Σ which is a π -graph and with boundary*

$$\partial\Sigma = \gamma \cup (\{p_1\} \times [0, +\infty)) \cup (\{p_2\} \times [0, +\infty)).$$

Moreover, Σ is nowhere vertical, is the unique minimal surface on $\mathbb{R}^2 \rtimes_A \mathbb{R}$ with such boundary and it is a Killing graph over the vertical domain $\Omega_\infty = \alpha \times [0, +\infty)$.

REMARK. Our construction works in some well studied spaces, for instance in the product space $\mathbb{H}^2 \times \mathbb{R}$, which is isometric and isomorphic to the semidirect product $\mathbb{R}^2 \rtimes_A \mathbb{R}$, when we choose

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$

In $\mathbb{H}^2 \times \mathbb{R}$, Scherk-like graphs have been already studied, and even more general results were obtained (for instance, in the work of Nelli and Rosenberg [NR] and in the work of Hauswirth, Rosenberg and Spruck [HRS]). However, the isometry between $\mathbb{R}^2 \rtimes_A \mathbb{R}$ and $\mathbb{H}^2 \times \mathbb{R}$ maps $\mathbb{R}^2 \rtimes_A \{0\}$ not to $\mathbb{H}^2 \times \{0\}$, as it could look at first sight, but to a horocylinder (that is, the product of a horocycle of $\mathbb{H}^2$ with $\mathbb{R}$), so the orientation of our graphs is not the usually studied in this space.

The proof of Theorem 4.1 is given in Section 4.2. If $\text{trace}(A) > 0$, when considering polygons P_c as in (40), there is a minimal π -graph Σ_c with boundary P_c . However, as the maximum principle does not hold, there is no reason for Σ_c be a Killing graph over Ω_c and we do not have the tools to ensure the continuity of the family Σ_c , which makes impossible to use geometric barriers. It becomes clear that, when $\text{trace}(A) \neq 0$, another sequence of surfaces Σ_c should be constructed, or other tools (such as stability of minimal π -graphs – a question that remains open) developed.

Our approach will be as follows: instead of considering minimal π -graphs over a domain on $\mathbb{R}^2 \rtimes_A \{0\}$, we will look to the problem *horizontally*, and consider an exhaustion of the *half-strip* $\Omega_\infty = \alpha \times [0, +\infty)$ by subdomains Ω_c on which is possible to find a family of minimal Killing graphs with prescribed boundary. Then, we use techniques from Killing graphs and elliptic partial differential equations to obtain the convergence of such family to another minimal Killing graph Σ_∞ . Then, we go back to the problem *vertically* (as the intermediate Killing graphs will also be π -graphs, by a result of Meeks, Mira, Pérez and Ros), and then we apply the geometric barriers used by A. Menezes to see that the surface Σ_∞ is, as claimed, a π -graph, nowhere vertical.

4.1. A good exhaustion of Ω_∞

Next proposition is crucial to the construction described above, as it gives the exhaustion of Ω_∞ by domains Ω_c where is possible to find minimal Killing graphs with prescribed boundary (see Figure 2).

PROPOSITION 4.2. *Let $\mathbb{R}^2 \rtimes_A \mathbb{R}$ be a semidirect product where $\text{trace}(A) \geq 0$. Then, there exists a constant $L_0 = L_0(A)$ depending uniquely on A such that for every two points $p_1, p_2 \in \mathbb{R}^2 \rtimes_A \{0\}$, if $\alpha = \overline{p_1 p_2}$ is the segment joining p_1 and p_2 and Ω_∞ is the vertical domain*

$$\Omega_\infty = \alpha \times [0, +\infty), \quad (41)$$

then, if $L = \text{length}(\alpha) < L_0$, Ω_∞ admits a continuous exhaustion $\{\Omega_c\}_{c>0}$ by domains Ω_c with boundary given by α , a graph over α , called α_c , and the two vertical segments joining the endpoints of α and α_c .

Moreover, such exhaustion is such that the Killing cylinder over $\partial\Omega_c$ with respect to the horizontal Killing field $Y_\theta = \sin(\theta)\partial_x + \cos(\theta)\partial_y$ has




Figure 2: The horizontal domain Δ and the exhaustion of Ω_∞ by subdomains Ω_c whose Killing cylinder (on the right) have mean curvature vector pointing inwards.

mean curvature vector pointing inwards, where θ is such that Y_θ is normal to Ω_∞ at $z = 0$.

PROOF. Let $p_1, p_2 \in \mathbb{R}^2 \rtimes_A \{0\}$ be any two points and, after a rotation on A as in (10) and a horizontal translation of $\mathbb{R}^2 \rtimes_A \mathbb{R}$, we may assume without loss of generality that $p_1 = (0, 0, 0)$ and $p_2 = (L, 0, 0)$ for some $L > 0$. We are going to show that if L is sufficiently small, then we can find the exhaustion as claimed.

In this setting, α is the segment $\alpha = \{(x, 0, 0) \mid 0 \leq x \leq L\}$ and Ω_∞ is the half-strip

$$\Omega_\infty = \{(x, 0, z) \in \mathbb{R}^2 \rtimes_A \mathbb{R} \mid 0 \leq x \leq L, z \geq 0\}, \quad (42)$$

transversal to the Killing field $Y = \partial_y$. Such assumptions will be kept until the end of the paper.

If $\text{trace}(A) = 0$, then the result is trivial (and without the need for an upper bound L_0) by taking α_c to be the translate of α to height c , $\alpha_c = \{(x, 0, c) \mid 0 \leq x \leq L\}$, as horizontal planes are minimal. Then, until the end of the proof we will treat the non-unimodular case and again we assume without loss of generality that $\text{trace}(A) = 2$, so A is a matrix as on (19). We will exhibit the curves α_c explicitly, then we prove they have the desired properties.

First, we treat the case where A is not diagonal and either $a^2 + bc \leq 0$ or $b \neq 0$: let λ, Λ the constants related with the matrix A via i . of Claim 1 and (25). Let

$$L_0 = \sqrt{\frac{\lambda}{2\Lambda}} \frac{\pi}{2}$$

and, for $L \leq L_0$, we let $f: [0, L] \rightarrow \mathbb{R}$ be

$$f(x) = \frac{1}{\Lambda} \ln \left(\frac{\cos \left(\sqrt{\frac{2\Lambda}{\lambda}} x \right)}{\cos \left(\sqrt{\frac{2\Lambda}{\lambda}} L \right)} \right). \quad (43)$$

Note that f is well defined, as $0 \leq x \leq L < L_0$ implies

$$\cos \left(\sqrt{\frac{2\Lambda}{\lambda}} x \right) \geq \cos \left(\sqrt{\frac{2\Lambda}{\lambda}} L \right) > 0,$$

so the quotient on (43) is larger than (or equal to) 1. In particular $f \geq 0$, with $f(x) = 0 \iff x = L$, and, for $c > 0$, we define $f_c = f + c$ and let $\alpha_c = \text{graph}(f_c) \subseteq \Omega_\infty$. Using such f_c , we define

$$\Omega_c = \{(x, 0, z) \in \mathbb{R}^2 \rtimes_A \mathbb{R} \mid 0 \leq x \leq L, 0 \leq z \leq f_c(x)\}, \quad (44)$$

and it follows that $\{\Omega_c\}_{c>0}$ is a continuous exhaustion of Ω_∞ . Next, we show that the Killing cylinder of the boundary of Ω_c with respect to ∂_y has mean curvature vector pointing inwards.

The ∂_y -Killing cylinder of $\partial\Omega_c$ has four smooth components (see Figure 2, right): one is a subdomain of a horizontal plane, so it has mean curvature 1 pointing upwards, two are contained on vertical planes, thus are minimal. The last component is the one corresponding to α_c , and it is a π -graph of the function $u_c(x, y) = f_c(x)$, hence (12) implies that its mean curvature, when oriented upwards, is

$$H = \frac{e^{4f_c}}{2W^3} \left[Q_{22}(f_c) f_c'' + G_1(f_c) (f_c')^2 + 2e^{-4f_c} \right]. \quad (45)$$

From the proof of Theorem 3.2, we obtain that $G_1/Q_{22} \leq \Lambda$. Moreover, Claim 1 implies that $Q_{22}(z) > \lambda e^{-2z}$, hence

$$H \leq \frac{e^{4f_c}}{2W^3} Q_{22}(f_c) \left[f_c'' + \Lambda (f_c')^2 + 2 \frac{e^{-2f_c}}{\lambda} \right], \quad (46)$$

whenever A is not diagonal and satisfies either $b \neq 0$ or $a^2 + bc \leq 0$. In particular, as $f_c \geq 0$ we have

$$H \leq \frac{e^{4f_c}}{2W^3} Q_{22}(f_c) \left[f_c'' + \Lambda (f_c')^2 + \frac{2}{\lambda} \right]. \quad (47)$$

Note that f was chosen in such a way it satisfies the ODE

$$f'' + \Lambda (f')^2 + \frac{2}{\lambda} = 0, \quad (48)$$

so, from (48) and (47), we obtain that $H \leq 0$, with respect to the upward orientation, hence the mean curvature vector of the Killing cylinder around α_c points downwards, as promised.




Figure 3: The surface Σ_c (on the right) is both a π -graph over Δ and a ∂_y -Killing graph over Ω_c , with $\partial\Sigma = \Gamma_c$ (the curve the left).

This finishes the proof when A is not diagonal and either $a^2 + bc \leq 0$ or $b \neq 0$. Now, we treat the simpler case of A being given by

$$A = \begin{pmatrix} 1+a & 0 \\ c & 1-a \end{pmatrix}. \quad (49)$$

It follows from (24) that $Q_{22}(z) = e^{2(a-1)z}$ and $G_1(z) = (3+a)e^{2(a-1)z}$, thus, the mean curvature of the π -graph of $u(x, y) = f(x)$ is

$$H = \frac{e^{2(a+1)f}}{2W^3} \left[f'' + (3+a)(f')^2 + 2e^{-2(1+a)f} \right],$$

and we can finish the proof similarly to the previous case. $\square$

4.2. Existence of Scherk-like graphs: Proof of Theorem 4.1

This section is to prove Theorem 4.1, done via an standard argument of convergence, with the difference that we look at the graphs sometimes vertically (as π -graphs), to have geometrically defined barriers, and sometimes horizontally (as ∂_y -Killing graphs), so we can use techniques of Killing graphs and elliptic partial differential equations.

PROOF (Proof of Theorem 4.1). Let $A \in M_2(\mathbb{R})$ be any matrix with $\text{trace}(A) \geq 0$ and let $L_0 > 0$ be the one given by Proposition 4.2. Let $p_1, p_2 \in \mathbb{R}^2 \rtimes_A \{0\}$ be such that $d(p_1, p_2) = L < L_0$ and without loss of generality assume $p_1 = (0, 0, 0)$ and $p_2 = (L, 0, 0)$.

Let $\alpha = \{(x, 0, 0) \mid 0 \leq x \leq L\}$ be the segment joining p_1 and p_2 and let $g: [0, L] \rightarrow \mathbb{R}$ be a convex, piecewise smooth function, with $g(0) = g(L) = 0$, meeting α on angles smaller than $\pi/2$ at 0 and L , that defines a piecewise smooth curve $\gamma \subseteq \mathbb{R}^2 \rtimes_A \{0\}$,

$$\gamma = \{(x, g(x), 0) \in \mathbb{R}^2 \rtimes_A \{0\} \mid 0 \leq x \leq L\},$$

with endpoints p_1, p_2 such that $\alpha \cup \gamma$ bounds a convex domain $\Delta \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ (as on Figure 3, left).

Let $\Omega_\infty = \alpha \times [0, +\infty)$ and, following the notation of Proposition 4.2, let, for each $c \geq 0$,

$$\Omega_c = \{(x, 0, z) \in \mathbb{R}^2 \rtimes_A \mathbb{R} \mid 0 \leq x \leq L, 0 \leq z \leq f_c(x)\}.$$

Let $\alpha_c = \{(x, 0, f_c(x)) \mid 0 \leq x \leq L\}$ be the graph of f_c , in such a way that its ∂_y -Killing cylinder

$$Cyl_{\partial_y}(\alpha_c) = \{(x, y, f_c(x)) \mid 0 \leq x \leq L, y \in \mathbb{R}\}$$

has mean curvature vector pointing downwards. We denote by

$$p_1(c) = (0, 0, f_c(0)), \quad p_2(c) = (L, 0, f_c(L))$$

the endpoints of α_c , and let, for $c \geq 0$, Γ_c be the simple closed curve in $\mathbb{R}^2 \rtimes_A \mathbb{R}$ given by (see Figure 3, left)

$$\Gamma_c = \gamma \cup \overline{p_1 p_1(c)} \cup \alpha_c \cup \overline{p_2 p_2(c)}. \quad (50)$$

CLAIM 3. The curve Γ_c as above bounds an *unique* minimal π -graph Σ_c over Δ , which is also a ∂_y -Killing graph over Ω_c .

Proof of Claim 3. First, we notice that Γ_c monotonically parametrizes $\partial\Delta$, then we can use Theorem 15.1 of [MMPR3] to obtain a minimal π -graph Σ_c with boundary $\partial\Sigma_c = \Gamma_c$.

First, we show that Σ_c is a ∂_y -Killing graph, on the sense that there is a function $g_c: \overline{\Omega}_c \rightarrow \mathbb{R}$, smooth up to the boundary, such that $R(g_c) = 0$, where R will stand for the elliptic operator of the mean curvature of minimal ∂_y -Killing graphs, and

$$\Sigma_c = Gr_{\partial_y}(g_c) = \{(x, g_c(x, z), z) \mid (x, 0, z) \in \Omega_c\}. \quad (51)$$

Note that, as Σ_c is a π -graph, there is a function $u_c: \Delta \rightarrow \mathbb{R}$ such that

$$\Sigma_c = \text{graph}(u_c) = \{(x, y, u_c(x, y)) \mid (x, y, 0) \in \Delta\}. \quad (52)$$

We claim Σ_c is contained in the ∂_y -Killing cylinder over Ω_c , so $0 \leq u_c(x, y) \leq f_c(x)$. Indeed, that $u > 0$ on the interior of Δ follows directly from the mean curvature comparison principle, so we show that $u_c(x, y) \leq f_c(x)$ for every $(x, y, 0) \in \Delta$. Arguing by contradiction, if there was an interior point $(x_0, y_0, 0) \in \Delta$ such that $u_c(x_0, y_0) > f_c(x_0)$, then we could consider the family $Cyl_{\partial_y}(\alpha_t)$, for $t > c$, and obtain a last contact point, interior for both Σ_c and $Cyl_{\partial_y}(\alpha_t)$, so the mean curvature of $Cyl_{\partial_y}(\alpha_t)$ would point upwards, a contradiction with Proposition 4.2 that proves that $u_c(x, y) \leq f_c(x)$ for every $(x, y, 0) \in \Delta$.

Let $q = (x, 0, z) \in \Omega_c$ be an interior point and consider $\mathcal{O}(q) = \{(x, y, z) \mid y \in \mathbb{R}\}$ the orbit of q with respect to the flux φ_t of the Killing field ∂_y . Note that $\mathcal{O}(q) \cap \Sigma_c$ is never empty for $q \in \Omega_c$, otherwise Σ_c would not be simply connected, hence it would not be a π -graph over Δ .

Moreover, the intersection $\mathcal{O}(q) \cap \Sigma_c$ cannot contain more than one point: if there were two points $q_i = \varphi_{t_i}(q) \in \Sigma_c$, with $t_1 < t_2$, then for $t_0 = t_2 - t_1 > 0$, $\varphi_{t_0}(\Sigma_c) \cap \Sigma_c \neq \emptyset$. Now, as $\varphi_t(\partial\Sigma_c) \cap \Sigma_c = \emptyset$ for all $t \neq 0$ by construction, we can consider the last contact point between $\varphi_t(\Sigma_c) \cap \Sigma_c$, and it will be interior for both Σ_c and $\varphi_t(\Sigma_c)$, a contradiction with the maximum principle.

This defines a function $g_c: \Omega_c \rightarrow \mathbb{R}$ satisfying the relation $(x, g_c(x, z), z) = \Sigma_c \cap \mathcal{O}(x, 0, z)$, thus Σ_c can be written as in (51). However, we still do not have the regularity of g_c . In order to prove that g_c is smooth, we begin by proving that the norm of $\text{grad}(g_c)$ is bounded in Ω_c .

Let $q \in \Omega_c$ be any interior point and consider a small open ball $B = B_{\Omega_c}(q, r) \subseteq \Omega_c$ such that $\text{Cyl}_{\partial_y}(\partial B)$ has mean curvature vector pointing inwards. Consider the following problem over $\overline{B}$:

$$\begin{cases} R(w) = 0, & \text{in } B \\ w|_{\partial B} = g_c|_{\partial B}, \end{cases} \quad (53)$$

where R is the mean curvature operator for ∂_y -Killing graphs. In other words, we are looking for a minimal ∂_y -Killing graph over a small ball on Ω_c that coincides with Σ_c on its boundary.

If $\Phi := g_c|_{\partial B}$ was of class $C^{2,\alpha}$, we could simply use the existence result due to M. Dajczer and J. H. de Lira, Theorem 1 of [DL]⁴ to obtain a solution to (53). However, at this point we can only guarantee that Φ is of class C^0 , so we need to use an approximation argument. Let $(\Phi_n^\pm)_{n \in \mathbb{N}} \subseteq C^{2,\alpha}(\partial B)$ be two sequences of $C^{2,\alpha}$ functions, converging to Φ and such that

$$\Phi_n^- \leq \Phi_{n+1}^- \leq \Phi \leq \Phi_{n+1}^+ \leq \Phi_n^+, \quad (54)$$

for every $n \in \mathbb{N}$. By Theorem 1 of [DL], there are functions $w_n^\pm \in C^{2,\alpha}(\overline{B})$ with minimal ∂_y -Killing graphs and such that $w_n^\pm|_{\partial B} = \Phi_n^\pm$. From (54) we obtain that the sequences $(w_n^\pm)_{n \in \mathbb{N}}$ are also monotone, $(w_n^-)_{n \in \mathbb{N}}$ is nondecreasing and $(w_n^+)_{n \in \mathbb{N}}$ is nonincreasing, both uniformly bounded. To obtain the convergence of the sequences $w_n^\pm$ to a solution of (53), we use some recent gradient estimates for Killing graphs obtained by J.-B. Casteras and J. Ripoll in [CR]:

THEOREM 4.3 (Theorem 4, [CR]). *Let M be a Riemannian manifold and let Y be a Killing field. Let Ω be a Killing domain in M and let $o \in \Omega$*

⁴We notice that the hypothesis on the Ricci curvature on [DL] is used uniquely to obtain an a priori estimate for the height of the graph, which is satisfied on our setting by the maximum principle.

and $r > 0$ such that the open geodesic ball $B_\Omega(o, r)$ is contained in Ω . Let $u \in C^3(B_\Omega(o, r))$ be a negative function whose Y -Killing graph has constant mean curvature H . Then there is a constant L depending only on $u(o)$, r , $|Y|$ and H such that $\|\text{grad}(u)(o)\| \leq L$.

All functions $w_n^\pm$ have uniform bounds on their C^0 norm, thus Theorem 4.3 above implies that there are uniform gradient estimates on compact subsets of B . This implies that both sequences will converge on the C^2 norm to a function $w \in C^2(B) \cap C^0(\overline{B})$, which is a solution of (53). Now, just use the flux of ∂_y and the same translation argument as before to obtain that w coincides with g_c in B , hence the gradient of g_c is bounded on interior points of Ω_c , as claimed.

Next, we use the relation $(x, g_c(x, z), z) = (x, y, u_c(x, y))$ to prove that g_c is actually smooth *up to the boundary*, with the unique exceptions of p_1 , p_2 , $p_1(c)$, $p_2(c)$ (where $\partial\Omega_c$ is not smooth), and the finite number of points where g is not differentiable. Note that u_c is smooth up to the boundary (except on the points where $\partial\Delta$ is not differentiable) and that the gradient of u_c is never horizontal on $\partial\Delta$, by the boundary maximum principle. Moreover, it follows from last argument that $\text{grad}(u_c)$ never vanishes on interior points of Δ , so g_c is also smooth up to the boundary, with the exceptions above.

This proves that any minimal π -graph S with $\partial S = \Gamma_c$ is a Killing graph. It follows easily that Σ_c is unique, proving Claim 3. $\diamond$

We proceed with the proof of Theorem 4.1, by noticing that the uniqueness of Σ_c , given by Claim 3, implies that the correspondence $c \mapsto g_c$ is continuous. Moreover, by its construction, we have that each g_c satisfies, on the boundary of Ω_c ,

$$g_c(0, z) = g_c(L, z) = 0, \quad g_c|_{\alpha_c} = 0, \quad g_c(x, 0) = g(x).$$

Again, as Σ_c is a π -graph over Δ , it is contained on the π -cylinder over Δ , and this can be translated to the horizontal setting as the inequality

$$0 \leq g_c(x, z) \leq g(x), \tag{55}$$

for every $(x, 0, z) \in \Omega_c$. Moreover, the usual argument using flux of ∂_y shows that the sequence g_c is monotonically increasing with c . In particular, as it is bounded, the sequence will converge pointwise for some function $g_\infty: \Omega_\infty \rightarrow \mathbb{R}$, such that $0 \leq g_\infty \leq g$. Next claim shows that the convergence is actually on the C^2 norm, so $\text{Gr}_{\partial_y}(g_\infty) = \Sigma_\infty$ is a minimal surface of $\mathbb{R}^2 \rtimes_A \mathbb{R}$.

CLAIM 4. When $c \rightarrow \infty$, the functions g_c converge on the C^2 norm to $g_\infty: \Omega_\infty \rightarrow \mathbb{R}$.

Proof of Claim 4. To prove this claim, we use the same argument of Claim 3, via gradient and height estimates for Killing graphs. Let $K \subseteq \overline{\Omega}_\infty$ be




Figure 4: $\Omega_\infty \subseteq P$ viewed horizontally: on each compact set $K \subseteq \overline{\Omega}_\infty$ there are uniform gradient estimates.

a compact set in $\overline{\Omega}_\infty$ with $C^{2,\alpha}$ boundary, as on Figure 4. As it holds $g_c(x, z) \leq g(x)$, it follows that the height of g_c is uniformly bounded on K , so we can use Theorem 4.3 to obtain a uniform bound for the norm of the gradient of g_c on interior points of K .

Note that (55), together with the assumption that the angle γ makes with α at p_1 and p_2 is less than $\pi/2$, implies that every g_c satisfies a uniform gradient estimate also along the boundary of K . As $g_c|_K \in C^{2,\alpha}(K)$ is smooth up to the boundary, this implies a uniform estimate for the gradient of g_c on K .

Now, just take an exhaustion of Ω_∞ by compact sets and, by using a standard argument, we obtain that a subsequence of (g_c) converges to g_∞ on the C^2 norm. In particular, as the sequence is monotone and converges pointwise, it follows that the convergence is smooth on the whole Ω_∞ . $\diamond$

From this claim we obtain that Σ_∞ is a minimal surface of $\mathbb{R}^2 \rtimes_A \mathbb{R}$, and that its boundary is

$$\partial\Sigma_\infty = \Gamma_\infty = \gamma \cup (\{p_1\} \times [0, \infty)) \cup (\{p_2\} \times [0, \infty)).$$

In order to finish the proof of Theorem 4.1, it remains to show that Σ_∞ is nowhere vertical and that it is unique. The uniqueness comes directly from the fact that it was built as a Killing graph, and that every other surface with such boundary is contained on the ∂_y -Killing cylinder over Ω_∞ .

To show that Σ_∞ is nowhere vertical, we go back to analyse the problem using π -graphs. First, if there was an interior point $p \in \Sigma_\infty$ such that $T_p\Sigma_\infty$ was vertical, Σ_∞ and $T_p\Sigma_\infty$ would be two minimal surfaces of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ tangent to each other at p . Then, there are at least two curves, meeting transversely at p on the intersection $T_p\Sigma_\infty \cap \Sigma_\infty$, so Σ_∞ cannot be a π -graph on a neighbourhood of p . Hence, it is a π -cylinder over some line segment⁵

⁵If $\beta \subseteq \mathbb{R}^2 \rtimes_A \{0\}$ is a smooth curve, the π -cylinder $\beta \times [0, \infty)$ is minimal if and only if β is a line segment: to see this, just use the foliation of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ by vertical planes which




Figure 5: The construction of the barrier $\tilde{R}$, by deforming ∂R over r_3 .

β contained on $\partial\Delta$. Secondly, if the point $p \in \partial\Sigma_\infty$ was a boundary point where $T_p\Sigma_\infty$ was vertical, then the boundary maximum principle would reach to the same conclusion. Next claim is to show that Σ_∞ meets $\pi^{-1}(\gamma)$ uniquely on γ , so $\Sigma_\infty \supseteq (\beta \times [0, \infty))$ is a contradiction.

CLAIM 5. $\Sigma_\infty \cap \pi^{-1}(\gamma) = \gamma$.

Proof of Claim 5. To prove this, we use the same barrier technique of A. Menezes, [Me]: Let γ_i be a smooth component of γ and let $p \in \gamma_i$ be any point. Consider L the vertical plane of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ containing the tangent line to γ_i at p (this is well defined even for $p \in \partial\gamma_i$, as γ_i is smooth). As γ is convex, this implies that Δ is contained on the same connected component of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ defined by L , and so does Σ_∞ .

Let, for $c \geq 0$, $u_c: \Delta \rightarrow \mathbb{R}$ be as in (52). and let $c_0 = \sup_\Delta u_0$. For $c_2 > c_1 > c_0$, consider a rectangle $R \subseteq L$ with boundary $\partial R = r_1 \cup r_2 \cup s_1 \cup s_2$ given by two parallel horizontal segments r_1 and r_2 and two vertical segments s_1 and s_2 , such that $s_1 \subseteq \{z = c_1\}$ and $s_2 \subseteq \{z = c_2\}$, that projects into $\mathbb{R}^2 \rtimes_A \{0\}$ in a compact segment $r \ni p$ with endpoints $q_1 = \pi(s_1)$ and $q_2 = \pi(s_2)$, contained on the same half-space determined by $\{y = 0\}$ (the vertical plane containing α) and with q_2 outside Δ , see Figure 5.

Let $q_3 \in \pi(R)$ be a point interior to the projection of R that is not in Δ . Then, $\tilde{q}_3 = \pi^{-1}(q_3) \cap r_2$ divides r_2 into two compact segments $r_3 \cup r_4$, r_3 projecting entirely outside Δ and with $p \in \pi(r_4)$.

are parallel to the vertical plane generated by the endpoints of β . It also follows from the more general formula $H(x, y, z) = k_g(x, y)e^{-z\text{trace}(A)}$, where $k_g(x, y)$ denotes the geodesic curvature of β on the point $(x, y, 0)$. The proof of this formula is a simple computation.

As L it is transversal to a (horizontal) Killing field, it is stable. In particular, it follows from the useful criteria due to D. Fischer-Colbrie and R. Schoen, Theorem 1 of [FS] (also proved on Proposition 1.32 of the book by Colding and Minicozzi, [CM]) that R is strictly stable, thus small perturbations of ∂R give rise to minimal surfaces with the perturbed boundary.

Change r_2 by making a parallel translation of r_3 on the direction of the half-space that contains Σ_∞ , whose projection still does not intersect Δ , joined by two small segments and denote such curve $\tilde{r}_3$, in such a way that $r_3 \cup \tilde{r}_3$ bounds a small rectangle on the horizontal plane $\{z = c_2\}$. We assume this perturbation is small in such way that its projection does not intersect Δ . Let $\tilde{R}$ be a minimal surface of $\mathbb{R}^2 \rtimes_A \mathbb{R}$ whose boundary is the perturbed rectangle $r_1 \cup \tilde{r}_3 \cup r_4 \cup s_1 \cup s_2$. Such surface is nowhere vertical and its contained in the convex hull of its boundary, in particular it is contained in $\{z \geq c_1\}$ and in the same half space that Σ_∞ with respect to L .

It is easy to see that $\pi(\tilde{R}) \cap \Delta \neq \emptyset$, otherwise $\tilde{R} \cap R$ would have a interior contact point. Moreover, $\tilde{R}$ is above u_0 on $\pi(\tilde{R}) \cap \Delta$, by the construction of $\tilde{R}$. Then, if $\Sigma_\infty \cap \pi^{-1}(\gamma_i) \neq \gamma_i$, we would have that $\Sigma_\infty \cap \tilde{R} \neq \emptyset$, thus, for some $\ell > 0$ there would be a first contact point between Σ_ℓ and $\tilde{R}$. As $\partial \Sigma_\ell$ does not intersect the convex hull of $\partial \tilde{R}$, it does not intersect $\tilde{R}$. Moreover, neither $\partial \tilde{R}$ can intersect Σ_ℓ , as this would imply such point would be in the plane L , so Σ_ℓ would have a vertical tangent plane. Then such contact point would be interior for both, reaching to a contradiction that proves the claim. $\diamond$

From Claim 5 and from the argument previously done, we obtain that Σ_∞ is a π -graph, nowhere vertical, which finishes the proof of Theorem 4.1. $\square$

References

- [CR] J.-B. Casteras and J. Ripoll, *On the asymptotic Plateau's problem for CMC hypersurfaces on rank 1 symmetric spaces of noncompact type*. to appear on Asian J. Math. preprint at <http://arxiv.org/abs/1403.1160>.
- [CM] T. H. Colding and W. P. Minicozzi II, *A course in minimal surfaces*. Number 121 in Graduate Studies in Mathematics. American Mathematical Society, 2011. ISBN-10: 0-8218-5323-6, ISBN-13: 978-0-8218-5323-8.
- [DL] M. Dajczer and J. H. de Lira, *Killing graphs with prescribed mean curvature and Riemannian submersions*. Ann. Inst. H. Poincaré Anal. Non Linéaire, **26**-3, (2009), p. 763-775.
- [FS] D. Fischer-Colbrie and R. Schoen, *The structure of complete stable minimal surfaces in 3-manifolds of nonnegative scalar curvature*. Comm. Pure Appl. Math. **33** (1980), no. 2, 199-211.

- [GT] D. Gilbarg and N. Trudinger, *Elliptic partial Differential Equations of Second Order*, 2nd edition. Springer - Verlag, (1983).
- [HRS] L. Hauswirth, H. Rosenberg and J. Spruck, *Infinite boundary value problems for constant mean curvature graphs in $\mathbb{H}^2 \times \mathbb{R}$ and $\mathbb{S}^2 \times \mathbb{R}$* , Amer. J. Math. **131** (2009), 195-226.
- [MMPR] W. H. Meeks III, P. Mira, J. Pérez and A. Ros, *Isoperimetric Domains of Large Volume in Homogeneous three-manifolds*. arXiv:1303.4222 [math.DG].
- [MMPR2] W. H. Meeks III, P. Mira, J. Pérez and A. Ros, *Constant mean curvature spheres in homogeneous three-spheres*. arXiv:1308.2612 [math.DG].
- [MMPR3] W. H. Meeks III, P. Mira, J. Pérez and A. Ros, *Constant mean curvature spheres in homogeneous three-manifolds*. Work in progress.
- [MP] W. H. Meeks III and J. Pérez, *Constant mean curvature surfaces in metric Lie groups*. Contemp. Math., **570** (2012) 25-110.
- [Me] A. Menezes, *Periodic minimal surfaces in semidirect products*. J. Aust. Math. Soc., **96** (2014) pp 127-144.
- [NR] B. Nelli and H. Rosenberg, *Minimal Surfaces in $\mathbb{H}^2 \times \mathbb{R}$* , Bull. Braz. Math. Soc., New Series **33** (2002), 263-292.
- [RST] H. Rosenberg, R. Souam and E. Toubiana, *General curvature estimates for stable H -surfaces in 3-manifolds and applications*. J. Differential Geom. **84** (2010), 623-648.

Álvaro Krüger Ramos
 Departamento de Matematica
 Univ. Federal do Rio Grande do Sul
 Av. Bento Gonçalves 9500
 91501-970 – Porto Alegre – RS – Brazil
 alvaro.ramos@ufrgs.br
 (supported by CNPq - Brasil)